



St Joseph's Catholic Primary School
Coast Road, Blackhall, Hartlepool, TS27 4HE
Tel/Fax: 0191 586 4308 Email: office@stjosephsblackhall.net
Website: www.st-josephsrc-ri.durham.sch.uk
Head Teacher: Mrs L Ashton Chair of Governors: Fr I Jackson



13th December 2023

Dear Parents/Carers,

Please find below government guidance for when children should be kept off school and when they shouldn't. As always if you do keep your child at home, it is important to contact school on the first day of absence to inform us of the reason.

Is my child too ill for school?

It can be tricky deciding whether or not to keep your child off school, when they're unwell. There are government guidelines for schools and nurseries about [managing specific infectious diseases at GOV.UK](#). These say when children should be kept off school and when they shouldn't. If you do keep your child at home, it's important to phone the school on the first day. Let them know that your child won't be in and give them the reason. If your child is well enough to go to school but has an infection that could be passed on, such as a cold sore or head lice, let their teacher know.

Other illnesses - Follow this advice for other illnesses:

Coughs and colds - It's fine to send your child to school with a minor [cough](#) or [common cold](#). But if they have a fever, keep them off school until the fever goes. Encourage your child to throw away any used tissues and to wash their hands regularly.

High temperature - If your child has a [high temperature](#), keep them off school until it goes away.

Chickenpox - If your child has [chickenpox](#), keep them off school until all the spots have crusted over. This is usually about 5 days after the spots first appeared.

Cold sores - There's no need to keep your child off school if they have a [cold sore](#). Encourage them not to touch the blister or kiss anyone while they have the cold sore, or to share things like cups and towels.

Conjunctivitis - You don't need to keep your child away from school if they have [conjunctivitis](#). Do get advice from your pharmacist. Encourage your child not to rub their eyes and to wash their hands regularly.

COVID-19 - If your child has mild symptoms, such as a runny nose, sore throat, or slight cough, and feels well enough, they can go to school. Your child should try to stay at home and avoid contact with other people if they have [symptoms of COVID-19](#) and they either:

- have a high temperature
- do not feel well enough to go to school or do their normal activities

What to do if your child has tested positive - Your child is no longer required to do a COVID-19 rapid lateral flow test if they have symptoms. But if your child has tested positive for COVID-19, they



St Joseph's Catholic Primary School
Coast Road, Blackhall, Hartlepool, TS27 4HE
Tel/Fax: 0191 586 4308 Email: office@stjosephsblackhall.net
Website: www.st-josephsrc-ri.durham.sch.uk
Head Teacher: Mrs L Ashton Chair of Governors: Fr I Jackson



should try to stay at home and avoid contact with other people for 3 days after the day they took the test.

Ear infection - If your child has an [ear infection](#) and a high temperature or severe earache, keep them off school until they're feeling better or their high temperature goes away.

Hand, foot and mouth disease - If your child has [hand, foot and mouth disease](#) but seems well enough to go to school, there's no need to keep them off. Encourage your child to throw away any used tissues straight away and to wash their hands regularly.

Head lice and nits - There's no need to keep your child off school once they have been treated for head lice. You can treat [head lice and nits](#) without seeing a GP.

Impetigo - If your child has [impetigo](#), they'll need treatment from a GP, often with antibiotics. Keep them off school until all the sores have crusted over and healed, or for 48 hours after they start antibiotic treatment. Encourage your child to wash their hands regularly and not to share things like towels and cups with other children at school.

Ringworm - If your child has [ringworm](#), see your pharmacist unless it's on their scalp, in which case you should see a GP. It's fine for your child to go to school once they have started treatment.

Scarlet fever - If your child has [scarlet fever](#), they'll need treatment with antibiotics from a GP. Otherwise they'll be infectious for 2 to 3 weeks. Your child can go back to school 24 hours after starting antibiotics.

Slapped cheek syndrome (fifth disease) - You don't need to keep your child off school if they have [slapped cheek syndrome](#) because, once the rash appears, they're no longer infectious. But let the school or teacher know if you think your child has slapped cheek syndrome.

Sore throat - You can still send your child to school if they have a [sore throat](#). But if they also have a high temperature, they should stay at home until it goes away. A sore throat and a high temperature can be symptoms of [tonsillitis](#).

Threadworms - You don't need to keep your child off school if they have [threadworms](#). Speak to your pharmacist, who can recommend a treatment.

Vomiting and diarrhoea - Children with [diarrhoea or vomiting](#) should stay away from school until they have not been sick or had diarrhoea for at least 2 days (48 hours).

Many thanks,

L Ashton
Executive Headteacher